HB 216 Great Bay Siltation Commission

September 12, 2007

Introduction to the siltation issue



What we know

- Channel morphology
- Turbidity
- Eelgrass
- Shellfish
- Dams


Channel morphology

- Federal channels in Squamscott, Lamprey, Cocheco, Bellamy and Piscataqua. Recent dredge projects on Squamscott and Cocheco. Condition reports are being completed or analyzed.
- Oyster "feasibility study for navigation."
 Significant deposition of silt in the upper third of the river.

Turbidity

 Have surface tidal or freshwaters shown a significant change in turbidity over time?

 TSS concentrations increased from an average value of 8.8 mg/L to 15.9 mg/L (an 81% increase) between 1976-1981 and 1999-2004 (NHEP, 2006)




• 1994 - 2004 TSS increased by approx. 20% at three major tributaries over a period when annual river flows went down.

 Sediment yield nearly doubled in that time period for the Oyster River.

 Together, the suspended solids from erosion and organic matter appear to be a significant factor that is limiting for light penetration to eelgrass and other underwater habitats


Eelgrass



Eelgrass

 New Hampshire Estuaries Project has documented a 17% loss in expanse of eelgrass since 1996.

- Approximately 1,500 acres have been lost since historic times.
- Eelgrass densities also appear to be decreasing.




Shellfish

 UNH project was smothered by sediment from May 2006 floods.

 Shellfish has declined for many years. It is at or near historic lows.

 Historically, oysters could filter all of Great Bay in less than 4 days. Today, it takes about 137 days.



Potential Shellfish restoration sites






Dams

 There are about 5,000 dams in New Hampshire!

- Each of the tributary rivers to Great Bay is blocked by one or more dams.
- Most of 1st or 2nd dams on the tributaries has been rebuilt or greatly modified in the last 70 years.

Dams around Great Bay



What we don't know

- Sources of sediments?
- Impacts of waves and wind?
- How much are human uses and ecological health being impacted?
- How much is human induced, how much is "natural"?
- How does sedimentation relate back to climate change?
- What are potential remedies?